



INTRODUCTION

This map displays part of the information requested by the City of Menlo Park to assist them in determining the location, quality and quantity of ground water. Information for the map was derived largely from confidential water well reports at the California Department of Water Resources and the City of Palo Alto. A few records were obtained from reports by Wood (1979), Hooper and others (1967), Sokol (1964), Poland (1970), and the California Water Commission (1950). The map by Hazlewood (1976) derived from the interpretation of seismic refraction profiles was used to determine the elevation of the bedrock surface in the area of salt evaporators north of Bayshore Freeway.

PROCEDURES

All of the water well information was organized by township, range and section. The location of each well was checked on published and unpublished maps or in the field. In many instances, the wells could not be accurately located and were omitted. Other wells are only approximately located within about 1,000 ft. Many more wells probably exist in the area of this map that have never been reported to State or local agencies.

The elevation of the bedrock surface in these wells that penetrated bedrock was derived by subtracting the depth to bedrock from the elevation of the well on the well log or on elevation estimated from the topographic map. These elevations are probably accurate within 10 ft, except for those in wells in Table 1 where the identification of the bedrock is queried. For these wells that did not penetrate bedrock, the elevation of the bottom of the hole was determined by subtracting from the total depth of the hole the elevation of the well at the surface.

Contours of the bedrock surface elevation were drawn by machine using a computerized algorithm which finds a surface of minimum total curvature, or the smoothest surface consistent with the data. This algorithm produces a contour map which makes the most neutral mathematical assumption, smoothness and flatness, in areas where no data are provided. The computer is from the Surface Cridding Library, prepared by the Software Development Group, Dynamic Graphics, Inc., Berkeley, California.

In unpublished topographic and well location map prepared in 1931 by someone at Stanford University, possibly J.J. Poland, above elevations for wells in the Palo Alto, Menlo Park, and Atherton areas that are as much as 15 ft higher than elevations on the 1961 U.S. Geological Survey 7.5' quadrangle. Poland (1969, 1971) determined that subsidence in the Atherton and Menlo Park areas from 1924 to 1961 was only about one foot, so nearly all of the elevation differences are probably related to errors in preparation of the different topographic maps. Subsidence was not taken into account in the preparation of the bedrock surface map because it is so small.

Faults that might affect the bedrock surface within this map area have been added from data largely independent of the wells. The Belmont Hill fault has been tentatively extended south into Redwood City from surface exposures mapped by Pampeyan (1981) and Brabb and Olson (1965) in the Belmont and San Carlos area approximately 3 miles northwest of the western border of this map. Another inferred fault in Palo Alto roughly parallel to Hanover Street could be a southerly extension of the Belmont Hill fault, but well data in between are not sufficient to link these features. The fault near Hanover Street is inferred from a northeast-facing escarpment about 20 ft high, and from the generally very deep elevation of the bedrock surface northwest of this escarpment and the occurrence of bedrock close to the surface southwest of the escarpment.

The Redwood City fault was named by Brabb and Hanna (1961) for an aeromagnetic anomaly extending northwestward approximately 16 miles from Palo Alto through the shoreline area of Redwood City. The fault is inferred from the linear character and steep gradient of the magnetic anomaly. Brabb and Hanna used a line to depict the fault at 1:125,000 scale, but at the 1:24,000 scale of this map, the line covers a zone more than 2 miles wide. The aeromagnetic anomalies and gravity measurements associated with the fault zone indicate that the rock is mainly serpentinite. In hazard has yet been established.

The Atherton and San Francisco faults are inferred from gravity measurements, and are discussed in another section of this report.

INTERPRETATION

Identification of the material and geologic formation from data provided by drillers is highly subjective. A 1927 log of a well at Sepolia High School in Redwood City, for example, indicates that the driller penetrated more than 200 ft of granite, blue sandstone, and black rock before encountering sandy clay, blue sand, and more granite and slate. These geologic relations and rock types are highly unlikely, judging from geologic maps of that area by Pampeyan (1970) and Brabb and Pampeyan (1981) and from data in nearby wells. Our interpretation is that the granite and black rock are probably different volcanic rocks within the Franciscan assemblage of Jurassic and Cretaceous age, and that the so-called slate represents shale within this assemblage. The sandy clay and blue sand could be shale and sandstone of the Franciscan assemblage that have been softened by tectonic shearing or weathering.

Information about the character of bedrock encountered in the principal wells used to construct the bedrock surface map is provided in Table 1. Note that the terms used by the drillers to describe the bedrock and our interpretation of the geologic unit involved are both provided.

We were not successful in identifying the Santa Clara Formation of late Pliocene and early Pleistocene age in any well. This unit has poorly consolidated sandstone, mudstone and conglomerate that is distinguishable from surficial deposits of Quaternary age in outcrop mainly by the amount of

deformation the unit has undergone. The term "bedrock" as used on this map, therefore, is applied to units stratigraphically beneath and older than the Santa Clara Formation.

Wells to bedrock north of a line extending north-southeast from outcrops of the Franciscan assemblage on Eagle Hill in Redwood City to the U.S. Geological Survey center in Menlo Park and beyond encountered mainly a hard shale that we interpret as the Franciscan assemblage. In the southern part of Redwood City, Atherton, and Menlo Park, the wells penetrated soft "banded" and interbedded sandstone that are here assigned to an unnamed sandstone, claystone and siltstone of early and middle Miocene age. This unit was referred to as an unnamed(?) sandstone on the map by Brabb and Pampeyan (1981). Some of the rocks could also be interpreted as those of the Franciscan assemblage. In the area of the central part of Stanford University southwest to the U.S. Veterans Hospital and beyond, the wells penetrated sandstone, shale, conglomerate and fossil shells that are probably from an unnamed formation of middle and late(?) Miocene age. For additional information about these geologic units, refer to the map by Brabb and Pampeyan (1981).

Spacing of wells to bedrock is not sufficient to determine which features outlined by the contours are related to tectonic erosion, errors in the data, or errors in interpreting the data. The abrupt change in elevation from bedrock at the surface along San Francisco Creek near Oak Knoll School to -537 ft is probably caused by movement along the Pulgas fault. The relative sense of movement is the northeast block down, but thrusting or lateral movement are also possible. The crowded, structurally-high areas near Chestnut Street in Redwood City and Palo Alto High School in Redwood City and Coyote Point in the northern part of San Mateo County. Some of the structurally low areas could be related to valleys carved by San Francisco Creek and other creeks thousands of years ago, or to tectonic doming. Alluvial deposits along the middle of the possible valleys would probably be thicker, coarser-grained, and more permeable.

ACKNOWLEDGMENTS

We are grateful to Philip Bellini, Donna Knifong, Robert Sikora and Nora Van Houten of the U.S. Geological Survey for helping collect well data in the Sacramento office of the California Department of Water Resources and to Betty Swanson of that organization for making the materials available. Billie, Patricia, Shirley, Van Houten, Betty Barred, Peter Decker, and David Walters kindly checked the location of many wells in the field. Fred Taylor, U.S. Geological Survey, collected additional information from David O'Leary, City of Palo Alto; Cheryl Jensen, Manager of Water Operation, City of San Mateo; and Richard Rose, Stanford University. Don Van Valen and Robert Banks of CUMR&S, San Francisco. We are also grateful to James to Niro, Medical Director, Stanford University, for loaning us the hospital area map, available information on water wells in the hospital area on Willow Road, Menlo Park, and to Walter Hensolt of Chin and Hensolt Engineers, Inc. for information on depth to bedrock throughout San Mateo County.

Thomas Poser, Richard Murrice, and John Klein, U.S. Geological Survey, provided information on wells drilled to bedrock at the U.S. Geological Survey Center in Menlo Park, and in the salt evaporator area near Dublin and Sausalito.

Brabb and Knifong are further acknowledged for digitizing the well information and for preparing preliminary copies of the bedrock surface map.

REFERENCES

- Brabb, E.E., and Hanna, W.F., 1961, Maps showing aeromagnetic anomalies, faults, earthquake epicenters, and igneous rocks in the southern San Francisco Bay region. U.S. Geological Survey Geophysical Map G-337, scale 1:125,000.
- Brabb, E.E., and Pampeyan, E.H., compilers, 1981, Geologic map of San Mateo County, California. U.S. Geological Survey Miscellaneous Investigations Series Map I-1257-F, scale 1:125,000.
- Brabb, E.E., and Olson, J.J., 1965, Map showing faults and earthquake epicenters in San Mateo County, California. U.S. Geological Survey Miscellaneous Investigations Series Map I-1257-F, scale 1:125,000.
- California State Water Resources Board, 1956, Santa Clara Valley investigations. California State Water Resources Board Bulletin, 7, 154 p.
- Dillehay, T.W., Jr., 1966, Geology of the Palo Alto quadrangle, Santa Clara and San Mateo Counties, California. California City, Mines and Geology Map Sheet 8, scale 1:62,500.
- Hazlewood, R.L., 1976, Contact map and interpretive cross sections showing depth and configuration of bedrock surface, south of San Francisco Bay region, California. U.S. Geological Survey Miscellaneous Field Studies Map MF-796, scale 1:62,500.
- Healy, J.R., and Taylor, R.L., 1979, Geology and engineering properties of the flatland deposits in flatland deposits of the San Francisco Bay region, California--their geology and engineering properties, and their importance to comprehensive planning. U.S. Geological Survey Professional Paper 943, p. 14-68.
- Hooper, C.W., Webb, J.L., and Newman, Robert, 1967, Effects of proposed salinity-control barriers in San Francisco Bay, California, upon groundwater resources. U.S. Geological Survey Miscellaneous Field Studies Map MF-575, scale 1:62,500.
- Pampeyan, E.H., 1970, Geologic map of the Palo Alto 7 1/2-minute quadrangle, San Mateo and Santa Clara Counties, California. U.S. Geological Survey Open-File Report 70-24, scale 1:24,000.
- Poland, J.J., 1961, Geology and former shoreline features of the San Mateo 7.5-minute quadrangle, San Mateo County, California. U.S. Geological Survey Open-File Report 61-39, scale 1:24,000.
- Poland, J.J., 1969, Land subsidence and aquifer-system compaction, Santa Clara Valley, California, USAI International Symposium on Land Subsidence, 1969, Sept. 1969.
- Poland, J.J., 1970, Ground water conditions in the Redwood City area, California, with particular reference to water supply for the Pacific Portland Cement Company. U.S. Geological Survey Open-File Report.
- Poland, J.J., 1971, Land subsidence in the Santa Clara Valley, Alameda, San Mateo, and Santa Clara Counties, California. U.S. Geological Survey Miscellaneous Field Studies Map MF-316, scale 1:125,000.
- Sokal, J.L., Menzies, R.L., and Lachar, R.L., 1966, Measurement of geothermal flux through poorly consolidated sediments: Earth and Planetary Science Letters 1, 1-10.
- Sokol, Daniel, 1964, The hydrogeology of the San Francisco Creek basin, San Mateo and Santa Clara Counties, California. Stanford University unpublished PhD thesis, 238 p.
- Wood, Percy, 1979, Sources of emergency water supplies in the San Mateo County, California. U.S. Geological Survey Open-File report 75-43, 25 p., map scale 1:125,000.